

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05393-B - Goat Anti-AIBZIP / CREB3L4, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CREB3L4, cAMP responsive element binding protein 3-like 4, AIBZIP, ATCE1, CREB3, CREB4, JAL, hJAL, CREB-4, androgen-induced basic leucine zipper protein, attaching to CRE-like 1, cAMP responsive element binding protein 1, cAMP-responsive element-binding protein 3-like protein 4, cAMP-responsive element-binding protein 4, cyclic AMP-responsive element-binding protein 4, tisp40, transcript induced in spermiogenesis protein 40

Official Symbol: CREB3L4

Accession Number(s): NP_570968.1; NP_001242909.1

Human GeneID(s): [148327](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_570968.1; NP_001242909.1). Reported variants represent identical protein: NP_001242908.1, NP_570968.1, NP_001242907.1 Reported variants represent identical protein: NP_001242909.1, NP_001242910.1

Immunogen

Peptide with sequence CPSGRIRSVLHADEM., from the C Terminus of the protein sequence according to NP_570968.1; NP_001242909.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 50kDa band observed in Human Placenta lysates (calculated MW of 43.4kDa according to NP_570968.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

Biotinylated EB05393 (1µg/ml) staining of Human Placenta lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.