

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06772 - Goat Anti-AKT2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AKT2, v-akt murine thymoma viral oncogene homolog 2, HGNC:392, PKBBETA, PRKBB, RAC-BETA, Murine thymoma viral (v-akt) homolog-2, rac protein

kinase beta

Official Symbol: AKT2

Accession Number(s): NP_001617.1; NP_001229956.1; NP_001317440.1

Human GeneID(s): 208

Important Comments: This antibody is expected to recognise all reported iasoforms

NP_001617.1; NP_001229956.1; NP_001317440.1.

Immunogen

Peptide with sequence C-RYDSLGLLELDQRT, from the C Terminus of the protein sequence according to NP_001617.1; NP_001229956.1; NP_001317440.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx. 58-60kDa band observed in Human Skeletal Muscle lysates and in Rat Testis lysates, and approx.60kDa in Human Testis lysates (calculated MW of 55.8kDa according to Human NP_001617.1 and 55.5kDa according to Rat NP_058789.1).

Recommended concentration: 0.5-1µg/ml. Primary incubation 1 hour at room temperature

Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog

EB06772 (1µg/ml) staining of Human Skeletal Muscle (A) and (0.75ug/ml) Human Testes (B) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB06772 (1μg/ml) staining of Rat Testes lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.