

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford

Oxfordshire

OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11714 - Goat Anti-Alpha-synuclein (NACP140) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: alpha-synuclein, NACP, non A-beta component of AD amyloid, PARK1, PARK4, PD1, SNCA, synuclein alpha-140, synuclein, alpha (non A4 component of amyloid precursor)

Official Symbol: SNCA

Accession Number(s): NP_000336.1

Human GeneID(s): 6622

Important Comments: This antibody is expected to recognize reported isoform NACP140 (NP_000336.1) only. Reported variants represent identical protein:

NP_000336.1, NP_001139527.1, NP_001139526.1

Immunogen

Peptide with sequence C-KDQLGKNEEGAPQE, from the internal region of the protein sequence according to NP_000336.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 17kDa band observed in Human Brain (Amygdala) lysates (calculated MW of 14.5kDa according to NP_000336.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.1-0.3μg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Brain (Cerebellum). Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Pig

Specific Reference

This antibody has been successfully used in WB and IF on Rat:

Muñoz P, Cardenas S, Huenchuguala S, Briceño A, Couve E, Paris I, Segura-Aguilar J. DT-Diaphorase Prevents Aminochrome-Induced Alpha-Synuclein Oligomer Formation and Neurotox Toxicol Sci. 2015 May:145(1):37-47.

PMID: 25634539

EB11714 (0.1μg/ml) staining of Human Amygdala lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.

EB11714 ($5\mu g/ml$) staining of paraffin embedded Human Cerebellum. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.