

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD

**Enquiries:** 

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05291 - Goat Anti-ASC / TMS1 Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** PYCARD, ASC, TMS1, PYD and CARD domain containing, CARD5, MGC10332, TMS, apoptosis-associated speck-like protein containing a CARD, caspase

recruitment domain protein 5, target of methylation-induced silencing-1, TMS-1

Official Symbol: PYCARD

Accession Number(s): NP\_037390.2; NP\_660183.1

Human GeneID(s): 29108

Important Comments: This antibody is expected to recognise both reported isoforms

(NP\_037390.2 and NP\_660183.1).

## Immunogen

Peptide with sequence C-RESQSYLVEDLERS, from the C Terminus of the protein sequence according to NP\_037390.2; NP\_660183.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 19kDa band observed in lysates of cell line HeLa (calculated MW of 20.0kDa according to NP\_660183.1). Recommended concentration: 0.2-0.6μg/ml. Primary incubation was 1 hour.

IHC: In paraffin embedded Human Liver shows staining of sinusoidal lining cells.

Recommended concentration: 4-6µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

EB05291 (0.2µg/ml) staining of HeLa lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB05291 (4µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.