

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08820 - Goat Anti-BMPR2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** BMPR2, bone morphogenetic protein receptor, type II (serine/threonine kinase), BMPR-II, BMPR3, BMR2, BRK-3, FLJ76945, PPH1, T-ALK, TRG10, bone morphogenetic protein receptor type II, primary pulmonary hypertension 1, serine/threonine kinase, transformation-related protein 10, type II activin receptor-like kinase

**Official Symbol:** BMPR2

**Accession Number(s):** NP\_001195.2

**Human GeneID(s):** [659](#)

### Immunogen

Peptide with sequence C-DLETNKLDPKVDK, from the internal of the protein sequence according to NP\_001195.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 110kDa band observed in Human Liver lysates (calculated MW of 115kDa according to NP\_001195.2). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow

### Specific Reference

**This antibody has been successfully used in the following paper:**

James W. Connell, Rachel J. Allison, Catherine E. Rodger, Guy Pearson, Eliska Zlamalova, Evan Reid

ESCRT-III-associated proteins and spastin inhibit protrudin-dependent polarised membrane traffic.

Cell Mol Life Sci. 2020 Jul;77(13):2641-2658.

PMID: 31587092

EB08820 (1µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.