

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

#### Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08821 - Goat Anti-CACNA1C Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** CACNA1C, calcium channel, voltage-dependent, L type, alpha 1C subunit, CACH2, CACN2, CACNL1A1, CCHL1A1, CaV1.2, MGC120730, TS, DHPR, alpha-1 subunit, calcium channel, L type, alpha 1 polypeptide, isoform 1, cardic muscle, calcium channel, cardic dihydropyridine-sensitive, alpha-1 subunit, voltage-dependent L-type calcium channel alpha 1C subunit, voltage-gated L-type calcium channel Cav1.2 alpha 1 subunit, splice variant 10\*, voltage-gated calcium channel alpha subunit Cav1.2 **Official Symbol:** CACNA1C

Accession Number(s): NP\_000710.5; NP\_955630.2; NP\_001161097.1; NP\_001123299.1; NP\_001123301.1; NP\_001161096.1; NP\_001123302.1; NP\_001123303.1 and more

## Human GeneID(s): 775

**Important Comments:** This antibody is expected to recognise all reported human isoforms.

## Immunogen

Peptide with sequence C-RARGRPSEEELQD, from the C Terminus of the protein sequence according to NP\_000710.5; NP\_955630.2; NP\_001161097.1; NP\_001123299.1; NP\_001123301.1; NP\_001161096.1; NP\_001123302.1; NP\_001123303.1 and more.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx. 200kDa band observed in Human Heart lysate (calculated MW of 240kDa according to NP\_000710.5). Recommended concentration: 1.0-3.0µg/ml.

## **Species Reactivity**

Tested: Human Expected from sequence similarity: Human EB08821 (1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.