

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06946 - Goat Anti-Cannabinoid Receptor 2 Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: cannabinoid receptor 2 (macrophage), HGNC:2160, CNR3, CB2, CX5, OTTHUMP00000044841, CNR2, RP11-4M23.1, testis-dominant CNR2 isoform CB2 Official Symbol: CNR2 Accession Number(s): NP_001832.1

Human GeneID(s): 1269

Immunogen

Peptide with sequence C-TETEADGKITPWPD, from the C Terminus of the protein sequence according to NP_001832.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 40kDa band observed in Human Brain (Hippocampus) lysates (calculated MW of 46.2kDa according to NP_001832.1). This molecular weight is routinely observed by other sources. Primary incubation was 1 hour. Recommended concentration: 1.5-4.5µg/ml. This antibody has been successfully used in WB on Human: Fügedi G et al. (2014) PMID: 25444073.

Species Reactivity

Tested: Human Expected from sequence similarity: Human

Specific Reference

This antibody has been successfully used in Western blot on Human: Fügedi G, Molnár M, Rigó J Jr, Schönléber J, Kovalszky I, Molvarec A. Increased placental expression of cannabinoid receptor 1 in preeclampsia: an observational study. BMC Pregnancy Childbirth. 2014 Dec 2;14:395 PMID: 25444073



EB06946 (1.5µg/ml) staining of Human Brain (Hippocampus) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.