

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05563 - Goat Anti-CAPE Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: SMC2, structural maintenance of chromosomes 2, PRO0324, CAP-E, CAPE, FLJ10093, SMC2L1, hCAP-E, OTTHUMP00000021818, OTTHUMP00000021819, SMC protein 2, SMC-2, SMC2 (structural maintenance of chromosomes 2, yeast)-like 1, SMC2 structural maintenance of chromosomes 2-like 1, XCAP-E homolog, chromosome-associated protein E, structural maintenance of chromosomes (SMC) family member, chromosome-associated protein E, structural maintenance of chromosomes protein 2 Official Symbol: SMC2

Accession Number(s): NP\_006435.2

Human GeneID(s): 10592

**Important Comments:** Reported variants NP\_006435.2, NP\_001036015.1 and NP\_001036016.1 represent identical protein

#### Immunogen

Peptide with sequence C-SKAKPPKGAHVEV, from the C Terminus of the protein sequence according to NP\_006435.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 140kDa band observed in lysates of cell line HeLa (calculated MW of 136kDa according to NP\_006435.2). Recommended concentration: 0.2-0.6µg/ml.

## **Species Reactivity**

Tested: Human Expected from sequence similarity: Human



EB05563 (0.2µg/ml) staining of HeLa lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.