

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06817 - Goat Anti-Caveolin 1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CAV1, caveolin 1, caveolae protein, 22kDa, HGNC:1527, CAV, MSTP085, VIP21, caveolae protein, 22-kD, caveolin 1, caveolin 1 caveolae protein, 22kD, caveolin 1, alpha isoform, caveolin 1, beta isoform, cell growth-inhibiting protein 32, BSCL3, CGL3, LCCNS

Official Symbol: CAV1

Accession Number(s): NP_001744.2; NP_001166366.1

Human GeneID(s): [857](#)

Non-Human GeneID(s): 12389 (mouse), 25404 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_001744.2; NP_001166366.1). Reported variants represent identical protein: NP_001166367.1, NP_001166368.1, NP_001166366.1. No cross-reactivity expected to Caveolin 2 and Caveolin 3

Immunogen

Peptide with sequence C-DELSEKQVYDAH, from the internal region of the protein sequence according to NP_001744.2; NP_001166366.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 22kDa band observed in lysates of cell lines A431 and A549 and in Human Heart lysates (calculated MW of 20.5kDa according to NP_001744.2).

Recommended concentration: 0.001-0.01µg/ml. Primary incubation 1 hour at room temperature. Preliminary testing was unsuccessful on Pig Heart and Spleen, and Mouse and Rat Lung for this particular batch.

IHC: Paraffin embedded Human Uterus. Recommended concentration: 2.5µg/ml.

Immunofluorescence: Strong expression of the protein seen in the Golgi and membranes of A431 cells and in the cytoplasm and Golgi of U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

EB06817 (0.003µg/ml) staining of A431 (A) and (0.001ug/ml) A549 cell lysate (35µg protein in RIPA buffer).
Detected by chemiluminescence.

EB06817 (0.003µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Detected by
chemiluminescence.

EB06817 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton.
Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Golgi and
membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml)
followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB06817 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton.
Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Golgi and
cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml)
followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB06817 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5%
Triton. Primary incubation overnight (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG
control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.

EB06817 (2.5µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer
pH 6, AP-staining.