



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07190 - Goat Anti-CDX2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CDX2, caudal type homeo box transcription factor 2, HGNC:1806, CDX-3, CDX3

**Official Symbol:** CDX2

**Accession Number(s):** NP\_001256.1

**Human GeneID(s):** [1045](#)

**Non-Human GeneID(s):** 12591 (mouse), 66019 (rat)

**Important Comments:** No cross-reactivity expected with CDX1 and 4

### Immunogen

Peptide with sequence EQLSPGGQRRNLCE, from the internal region of the protein sequence according to NP\_001256.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** Approx 35kDa band observed in Human Colon lysates (calculated MW of 33.5kDa according to NP\_001256.1). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Colon. Recommended concentration: 5µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog

EB07190 (1µg/ml) staining of Human Colon lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB07190 (5µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.