

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09314 - Goat Anti-CENTG1 / PIKE Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CENTG1, centaurin, gamma 1, AGAP2, FLJ16430, GGAP2, KIAA0167, PIKE, Arf GAP with GTP-binding protein-like, ANK repeat and PH domains 2, GTP-binding and GTPase activating protein 2, centaurin gamma1, phosphoinositide 3-kinase enhancer

**Official Symbol:** AGAP2

**Accession Number(s):** NP\_055585.1

**Human GeneID(s):** [116986](#)

**Important Comments:** This antibody is expected to recognize isoform PIKE-S (NP\_055585.1).

### Immunogen

Peptide with sequence C-QASLDSIREAVINSQ, from the internal region of the protein sequence according to NP\_055585.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 100kDa band observed in Human Brain (Amygdala and Cerebral Cortex) lysates (calculated MW of 90.5kDa according to NP\_055585.1). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Thyroid and Kidney. Recommended concentration: 3.75µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human

EB09314 (0.3µg/ml) staining of Human Amygdala lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB09314 (3.75µg/ml) staining of paraffin embedded Human Thyroid. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB09314 (3.75µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.