

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09305 - Goat Anti-CLTC Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CLTC, clathrin, heavy chain (Hc), CHC17, CLH-17, CLTCL2, Hc, KIAA0034, clathrin heavy chain 1, clathrin, heavy polypeptide (Hc), clathrin, heavy polypeptide-like 2

**Official Symbol:** CLTC

**Accession Number(s):** NP\_004850.1; NP\_001275582.1

**Human GeneID(s):** [1213](#)

### Immunogen

Peptide with sequence C-ESLRKEEEQATETQ, from the internal region (near C Terminus) of the protein sequence according to NP\_004850.1; NP\_001275582.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**IHC:** In paraffin embedded Human Adrenal Gland shows strong vesicular staining in the zona glomerulosa. Paraffin embedded Human Kidney. Recommended concentration: 3.75µg/ml.

**Immunofluorescence:** This antibody has been successfully used in IF on Human: Marinval N et al. (2016) PMID: 27763505.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow

### Specific Reference

**This antibody has been successfully used in IF on Human:**

Marinval N, Saboural P, Haddad O, Maire M, Bassand K, Geinguenaud F, Djaker N, Ben Akrou K, Lamy de la Chapelle M, Robert R, Oudar O, Guyot E, Laguillier-Morizot C, Sutton A, Chauvierre C, Chaubet F, Charnaux N, Hlawaty H  
Identification of a Pro-Angiogenic Potential and Cellular Uptake Mechanism of a LMW Highly Sulfated Fraction of Fucoidan from Ascophyllum nodosum  
Mar Drugs. 2016 Oct 17;14(10)  
PMID: 27763505

EB09305 (3.75µg/ml) staining of paraffin embedded Human Adrenal Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB09305 (3.75µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.