

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06532 - Goat Anti-CREM Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CREM, ICER, MGC17881, MGC41893, MGC111110, hCREM-2, cAMP responsive element modulator, hCREM 2alpha-b protein, hCREM 2beta-a protein, inducible cAMP early repressor ICER, OTTHUMP00000019442,

OTTHUMP00000019443, cAMP response element modulator; hCREM 2alpha-b protein **Official Symbol:** CREM

Accession Number(s): NP_853549.1; NP_874387.1; NP_874388.1; NP_874393.1; NP_874394.1; NP_877570.1; NP_877571.1; NP_877572.1; NP_877573.1; NP_878270.1; NP_898883.1

Human GeneID(s): 1390

Important Comments: This antibody is expected to recognise the following protein isoforms of human CREM protein: a (NP_853549.1); e (NP_874387.1); f (NP_874388.1); k (NP_874393.1); I (NP_874394.1); m (NP_877570.1); n (NP_877571.1); o (NP_877572.1); p (NP_877573.1); q (NP_878270.1); v (NP_898883.1). This antibody is not expected to cross-react with CREB protein.

Immunogen

Peptide with sequence LETLKDICSPKTDY, from the C Terminus of the protein sequence according to NP_853549.1; NP_874387.1; NP_874388.1; NP_874393.1; NP_874394.1; NP_877570.1; NP_877571.1; NP_877572.1; NP_877573.1; NP_878270.1; NP_898883.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 35kDa band observed in Human Heart lysates (calculated MW of 32.5kDa according to NP_853549.1). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow

EB06532 staining (1µg/ml) of Human Heart lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.