

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB12710 - Goat Anti-CSNK2B Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CSNK2B, casein kinase 2, beta polypeptide, CK2B, CK2N, CSK2B, G5A, CK II beta, Casein kinase II beta subunit, alternative name: G5a, phosvitin, casein

kinase II subunit beta, phosvitin, protein G5a

Official Symbol: CSNK2B

Accession Number(s): NP_001311.3; NP_001269314.1

Human GeneID(s): 1460

Non-Human GenelD(s): 13001 (mouse), 81650 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms

(NP_001311.3; NP_001269314.1).

Immunogen

Peptide with sequence C-TGLNEQVPHYRQ, from the internal region of the protein sequence according to NP_001311.3; NP_001269314.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 26kDa band observed in lysates of cell lines NIH3T3, HeLa, K562, MCF7 and HepG2 (calculated MW of 24.9kDa according to NP_001311.3). Approx 25kDa band observed in Mouse Testis, 26kDa in Rat Testis and 28kDa in Pig Testis lysates (calculated MW of 24.9kDa according to Mouse NP_001290374.1, Rat NP_001030315.1 and pig NP_001138849.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.3-1μg/ml.

Species Reactivity

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB12710 (1µg/ml) staining of NIH3T3 (A), HeLa (B), K562 (C), MCF7 (D) and HepG2 (E) lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB12710 (0.3µg/ml) staining of Mouse (A), Rat (B) and Pig (C) Testis lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.