

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com

Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB12525 - Goat Anti-CXCR1 (aa26-38) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CXCR1, chemokine (C-X-C motif) receptor 1, C-C, C-C-CKR-1, CD128, CD181, CDw128a, CKR-1, CMKAR1, IL8R1, IL8RA, IL8RBA, C-X-C chemokine receptor type 1, CXC-R1, CXCR-1, IL-8 receptor type 1, IL-8R A, high affinity interleukin-8 receptor A, interleukin 8 re Official Symbol: CXCR1 Accession Number(s): NP_000625.1 Human GeneID(s): <u>3577</u>

Immunogen

Peptide with sequence DYSPCMLETETLN, from the internal region (near N terminus) of the protein sequence according to NP_000625.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000. **Western blot:** Approx 39kDa band observed in Human Breast cancer lysates (calculated MW of 39.8kDa according to NP_000625.1). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human EB12525 (0.3µg/ml) staining of Human Breast cancer lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.