

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

#### Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07126 - Goat Anti-DEAD-box protein 6 Antibody 🏹

Size: 100µg specific antibody in 200µl

## **Target Protein**

Principal Names: DEAD-box protien 6, DEAD (Asp-Glu-Ala-Asp) box polypeptide 6, DDX6, HLR2, P54, RCK, DEAD box-6, DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 6 (RNA helicase, 54kD) Official Symbol: DDX6 Accession Number(s): NP\_004388.1 Human GenelD(s): <u>1656</u>

#### Immunogen

Peptide with sequence C-AEYHSEPVEDEKP, from the C Terminus of the protein sequence according to NP\_004388.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

**Peptide ELISA:** antibody detection limit dilution 1:128000. **Western blot:** Approx 50kDa band observed in lysates of cell lines Daudi, Jurkat and HepG2 (calculated MW of 53.2kDa according to NP\_004388.1). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour. **IHC:** Paraffin embedded Human Brain (Cortex) and Thymus. Recommended concentration: 5µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

EB07126 (0.3µg/ml) staining of Daudi (A), Jurkat (B) and HepG2 (C) lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB07126 (5µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB07126 (5µg/ml) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.