

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10981 - Goat Anti-FUT2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FUT2, fucosyltransferase 2 (secretor status included), B12QTL1, SE, SEC2, Se2, sej, GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase 2, alpha (1,2) fucosyltransferase, alpha(1,2)FT2, galactoside 2-alpha-L-fucosyltransferase 2 Official Symbol: FUT2 Accession Number(s): NP_000502.4 Human GenelD(s): <u>2524</u> Important Comments: Reported variants represent identical protein: NP_001091107.1, NP_000502.4

Immunogen

Peptide with sequence C-HVQQRLAKIQAMWE, from the internal region (near N Terminus) of the protein sequence according to NP_000502.4.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 37kDa band observed in Human Liver lysates (calculated MW of 39.0kDa according to NP_000502.4. Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Placenta. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human EB10981 (1µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB10981 (5µg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.