

# **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05153 - Goat Anti-FYN Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** FYN, SLK, SYN, MGC45350, FYN oncogene related to SRC, FGR, YES, src-like kinase, c-syn protooncogene, tyrosine kinase p59fyn(T), src/yes-related novel gene, OKT3-induced calcium influx regulator, proto-oncogene tyrosine-protein kinase fyn, RP1-66H14.1, OTTHUMP00000017917, protein-tyrosine kinase fyn, src-like kinase

Official Symbol: FYN Accession Number(s): NP\_002028.1; NP\_694592.1; NP\_694593.1 Human GenelD(s): 2534 Non-Human GenelD(s): 14360 (mouse), 25150 (rat) Important Comments: This antibody is expected to recognise all three reported isoforms (as represented by NP\_002028.1; NP\_694592.1; NP\_694593.1).

## Immunogen

Peptide with sequence VQCKDKEATKLTE, from the N Terminus of the protein sequence according to NP\_002028.1; NP\_694592.1; NP\_694593.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 55kDa band observed in lysates of cell lines A431 and Jurkat (calculated MW of 54.5kDa according to NP\_694593.1). Recommended for use at  $0.3-1\mu g/ml$ .

# **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow EB05153 (0.3µg/ml) staining of A431 (A) and Jurkat lysate (B) (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.