



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05570 - Goat Anti-GATA3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** GATA3, GATA binding protein 3, HDR, MGC2346, MGC5199, MGC5445, GATA-binding protein 3, enhancer-binding protein GATA3

**Official Symbol:** GATA3

**Accession Number(s):** NP\_001002295.1; NP\_002042.1

**Human GeneID(s):** [2625](#)

**Non-Human GeneID(s):** 14462 (mouse)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_001002295.1 and NP\_002042.1).

### Immunogen

Peptide with sequence EVTADQPRWVSHH-C, from the N Terminus of the protein sequence according to NP\_001002295.1; NP\_002042.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 46kDa band observed in Human Tonsil and in Mouse Kidney lysates. (calculated MW of 47.9kDa according to NP\_002042.1 and Mouse NP\_032117.1). Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human, Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow

EB05570 (0.5µg/ml) staining of Human Tonsil (A) and Mouse Kidney (B) lysate (35µg protein in RIPA buffer).  
Primary incubation was 1 hour. Detected by chemiluminescence