



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10742-T - Goat Anti-GNAQ / ALPHA-q (aa162-175) Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: GNAQ, guanine nucleotide binding protein (G protein), q polypeptide, G-ALPHA-q, GAQ

Official Symbol: GNAQ

Accession Number(s): NP_002063.2

Human GeneID(s): [2776](#)

Non-Human GeneID(s): 14682 (mouse), 81666 (rat)

Immunogen

Peptide with sequence C-NDLDRVADPAYLPT, from the internal region of the protein sequence according to NP_002063.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 40kDa band observed in Human, Mouse and Rat Testis lysates, in Pig Colon and Spleen lysates and in lysates of Human Peripheral Blood Mononucleocytes (calculated MW of 42.1kDa according to Human NP_002063.2 and to Rat NP_112298.1). Recommended concentration: 0.1-0.3µg/ml.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB10742 (0.1µg/ml) staining of Mouse (A) and Rat (B) Testis lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB10742 (0.1µg/ml) staining of Peripheral Blood Lymphocytes lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB10742 (0.1µg/ml) staining of Pig Colon (A) and Pig Spleen (B) lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB10742 (5µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.