

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09623 - Goat Anti-HMGI-C / HMGA2 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** HMGA2, high mobility group AT-hook 2, BABL, HMGI-C, HMGIC, LIPO, STQTL9, High-mobility group protein HMGI-C, high-mobility group (nonhistone

chromosomal) protein isoform I-C

Official Symbol: HMGA2

Accession Number(s): NP\_003474.1; NP\_003475.1

Human GeneID(s): 8091

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_003474.1; NP\_003475.1) and is not expected to cross-react with HMGA1.

# Immunogen

Peptide with sequence C-KAAQKKAEATGEK, from the internal region of the protein sequence according to NP\_003474.1; NP\_003475.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 18kDa band observed in Human Heart lysates (calculated MW of 11.8kDa according to NP\_003474.1 and 11.5kDa according to NP\_003475.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Di Cello et al, Mol Cancer Res. 2008 May;6(5):743-50.; PMID: 18505920). Recommended concentration: 1-3μg/ml.

rtecommended concentration. 1-5μξ

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Dog

EB09623 ( $1\mu g/ml$ ) staining of Human Heart lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.