

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09130 - Goat Anti-KCNJ1 / ROMK Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** KCNJ1, potassium inwardly-rectifying channel, subfamily J, member 1, KIR1.1, ROMK, ROMK1, ATP-regulated potassium channel ROM-K, ATP-sensitive inward rectifier potassium channel 1, OTTHUMP00000045938, inwardly rectifying K+ channel,

potassium inwardly-rectifying channel J1

Official Symbol: KCNJ1

Accession Number(s): NP\_000211.1; NP\_722448.1

Human GeneID(s): 3758

Non-Human GeneID(s): 56379 (mouse), 24521 (rat)

**Important Comments:** This antibody is expected to recognize reported isoforms NP\_722449.2 and NP\_000211. The following reported variants represent identical protein:

NP\_722451.1, NP\_722449.2, NP\_722450.1, NP\_722448.1.

### **Immunogen**

Peptide with sequence C-DQININFVVDAGNEN, from the internal region of the protein sequence according to NP\_000211.1; NP\_722448.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx. 45kDa band observed in Human Kidney lysates (calculated MW of 44.8kDa according to NP\_000211.1). Recommended concentration: 1-3μg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

EB09130 (1μg/ml) staining of Human Kidney lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.