

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09445 - Goat Anti-LMP7 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PSMB8, proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional peptidase 7), D6S216, D6S216E, LMP7, MGC1491, PSMB5i, RING10, beta5i, OTTHUMP00000029420, OTTHUMP00000029421, OTTHUMP00000062981, low molecular weight protein 7, macropain subunit C13, multicatalytic endopeptidase complex subunit C13, protease component C13, proteasome (prosome, macropain) subunit beta type 8 (large multifunctional protease 7), proteasome beta 8 subunit, proteasome catalytic subunit 3i, proteasome subunit Y2, proteasome subunit beta 5i, proteasome-related gene 7

**Official Symbol:** PSMB8

**Accession Number(s):** NP\_004150.1; NP\_683720.2

**Human GeneID(s):** [5696](#)

**Non-Human GeneID(s):** 16913 (mouse), 24968 (rat)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_004150.1; NP\_683720.2).

### Immunogen

Peptide with sequence C-DVSDLLHQYREANQ, from the C Terminus of the protein sequence according to NP\_004150.1; NP\_683720.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 22kDa band observed in Human Liver, Lung and Lymph Node lysates (calculated MW of 29.8kDa according to precursor NP\_004150.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Schwarz et al, J Immunol. 2000 Jul 15;165(2):768-78. PMID: 10878350. Recommended concentration: 0.01-0.03µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Spleen. Recommended concentration: 3.75µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow



EB09445 (0.01µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB09445 (3.75µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.