

### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09756 - Goat Anti-MLX Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** MLX, MAX-like protein X, MAD7, MXD7, TCFL4, bHLHd13, BigMax protein, MAX-like bHLHZIP protein, transcription factor-like 4, transcription factor-like

protein 4

Official Symbol: MLX

Accession Number(s): NP\_937848.1; NP\_937847.1; NP\_733752.1

Human GeneID(s): 6945

Non-Human GenelD(s): 21428 (mouse), 360631 (rat)

Important Comments: This antibody is expected to recognize all three reported isoforms

(NP\_937848.1; NP\_937847.1; NP\_733752.1).

## **Immunogen**

Peptide with sequence C-QIVKAHQDNPHEGED, from the internal region of the protein sequence according to NP\_937848.1; NP\_937847.1; NP\_733752.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Preliminary experiments in Human, Mouse and Rat Liver lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml).

IHC: In paraffin embedded Human Placenta shows nuclear stsaining of trophoblasts.

Recommended concentration: 2.5-3.8µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

EB09756 (3.8μg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.