

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07039 - Goat Anti-MURF1 / TRIM63 (internal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TRIM63, tripartite motif-containing 63, HGNC:16007, FLJ32380, IRF, MURF1, MURF2, RNF28, SMRZ, iris ring finger protein, muscle specific ring finger protein 1, muscle specific ring finger protein 2, ring finger protein 28, striated muscle RING zinc finger protein

Official Symbol: TRIM63

Accession Number(s): NP_115977.2

Human GeneID(s): [84676](#)

Immunogen

Peptide with sequence C-QQYQEQLDKSTKLVE, from the internal region of the protein sequence according to NP_115977.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary experiments gave an approx 75kDa band in Human Heart and Skeletal Muscle lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 40.2kDa according to NP_115977.2. The 75kDa band was successfully blocked by incubation with the immunizing peptide.

IHC: Paraffin embedded Human Skeletal Muscle. Recommended concentration: 1.25µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

EB07039 (1.25µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.