

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05882 - Goat Anti-MURF2 / TRIM55 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** TRIM55, RNF29, MURF-2, ring finger protein 29, for muscle specific ring finger 2, tripartite motif-containing 55, muscle specific ring finger 2

**Official Symbol:** TRIM55

**Accession Number(s):** NP\_908973.1; NP\_908974.1; NP\_908975.1

**Human GeneID(s):** [84675](#)

**Non-Human GeneID(s):** 381485 (mouse), 365751 (rat)

**Important Comments:** This antibody is expected to recognize isoforms 1, 3 and 4 (NP\_908973.1, NP\_908974.1 and NP\_908975.1 resp.).

### Immunogen

Peptide with sequence C-HIFSFSWLNSLNE, from the C Terminus of the protein sequence according to NP\_908973.1; NP\_908974.1; NP\_908975.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 58kDa band observed in lysates of cell line HeLa (calculated MW of 60.5kDa according to NP\_908973.1). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature

**IHC:** Paraffin embedded Human Heart. Recommended concentration: 10µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the Golgi and cytoplasm of U2OS cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig

EB05882 (2µg/ml) staining of HeLa cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB05882 (10µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB05882 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Golgi and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB05882 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.