

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09020 - Goat Anti-MYH9 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: myosin, heavy chain 9, non-muscle, RP1-68O2.1, DFNA17, EPSTS, FTNS, MGC104539, MHA, NMHC-II-A, NMMHCA, MYH9 variant protein, OTTHUMP00000028706, cellular myosin heavy chain, type A, myosin, heavy polypeptide 9, non-muscle, non-muscle myosin heavy chain, non-muscle myosin heavy polypeptide 9, nonmuscle myosin heavy chain II-A Official Symbol: MYH9 Accession Number(s): NP_002464.1 Human GenelD(s): 4627

Immunogen

Peptide with sequence C-DQINTDLNLERSH, from the internal region of the protein sequence according to NP_002464.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 250kDa band observed in lysates of Human Peripheral Blood Mononucleocytes and of NIH3T3 (calculated MW of 227kDa according to Human NP_002464.1 and to Mouse NP_071855.2). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human, Mouse Expected from sequence similarity: Human, Mouse, Rat, Dog

Specific Reference

This antibody has been successfully used in Western blot on Human: Harms RZ, Borengasser K, Kumar V, Sarvetnick N Anti-human Interleukin (IL)-4 Clone 8D4-8 Cross-Reacts With Myosin-9 Associated With Apoptotic Cells and Should Not Be Used for Flow Cytometry Applications Querying IL-4 Expression. Front Cell Dev Biol. 2019 Apr 9;7:46. PMID: 31024909 EB09020 (0.3µg/ml) staining of Human peripheral blood lymphocytes lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB09020 (1µg/ml) staining of NIH3T3 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.