

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10519 - Goat Anti-NEK7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NIMA (never in mitosis gene a)-related kinase 7, NIMA-related kinase

7, NEK7

Official Symbol: NEK7

Accession Number(s): NP_598001.1

Human GenelD(s): 140609

Non-Human GenelD(s): 59125 (mouse), 360850 (rat)

Immunogen

Peptide with sequence C-NTLANFRIEKK, from the internal region of the protein sequence

according to NP_598001.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 35kDa band observed in lysates of cell line HeLa (calculated MW of 34.6kDa according to NP_598001.1). Recommended concentration: 0.05-0.2µg/ml.

Primary incubation was 1 hour.

IHC: Paraffin embedded Human Thyroid. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

EB10519 (0.05μg/ml) staining of HeLa lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.

EB10519 (3.75 μ g/ml) staining of paraffin embedded Human Thyroid. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.