



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10571 - Goat Anti-NPAS4 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NPAS4, neuronal PAS domain protein 4, Le-PAS, NXF, PASD10, bHLHe79, HLH-PAS transcription factor NXF, PAS domain-containing protein 10, class E basic helix-loop-helix protein 79, neuronal PAS4

**Official Symbol:** NPAS4

**Accession Number(s):** NP\_849195.2

**Human GeneID(s):** [266743](#)

**Non-Human GeneID(s):** 225872 (mouse), 266734 (rat)

### Immunogen

Peptide with sequence CRFNTSKSLRRQS, from the internal region of the protein sequence according to NP\_849195.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** Approx 90kDa band observed in Human Brain (Frontal Cortex) lysates and in Mouse and Rat Brain lysates (calculated MW of 87.1kDa according to NP\_849195.2).

Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human, Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow

EB10571 (0.3µg/ml) staining of Human Frontal Cortex lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB10571 (1µg/ml) staining of Mouse (A) and Rat (B) Brain lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.