

#### **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

## EB05061 - Goat Anti-P27KIP1 / CDKN1B (C Terminus) Antibody

Size: 100µg specific antibody in 200µl

# Target Protein



Principal Names: CDKN1B, P27KIP1, cyclin-dependent kinase inhibitor 1B (p27, Kip1), KIP1, CDKN4, cyclin-dependent kinase inhibitor 1B, p27(KIP1), KIP1, p27, MEN1B, MEN4, P27KIP1 Official Symbol: CDKN1B Accession Number(s): NP\_004055.1 Human GenelD(s): 1027 Non-Human GenelD(s): 12576 (mouse), 83571 (rat)

#### Immunogen

Peptide with sequence C-EQTPKKPGLRRRQT, from the C Terminus of the protein sequence according to NP\_004055.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 26-27kDa band observed in Human Breast and Prostate lysates (calculated MW of 22.1kDa according to NP\_004055.1). The observed molecular weight corresponds to earlier findings in literature (Xia et al, Proc Natl Acad Sci U S A. 2005 Sep 27;102(39):14028-33; PMID: 16169901). Recommended concentration: 0.5-1.5µg/ml.

**Immunoprecipitation:**This antibody was deemed fit for IP under native conditions (observations from anonymous customer).

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Cow, Dog



EB05061 (0.5µg/ml) staining of Human Breast lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.