



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10589 - Goat Anti-p47phox (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Ncf1, neutrophil cytosolic factor 1, NOXO2, Ncf-1, p47<phox>, p47phox, 47 kDa neutrophil oxidase factor, NADPH oxidase subunit (47 kD), NADPH oxidase subunit (47kDa), OTTMUSP00000022112, neutrophil NADPH oxidase factor 1, neutrophil cytosol facto

Official Symbol: Ncf1

Accession Number(s): NP_035006.3

Human GeneID(s): [653361](#)

Non-Human GeneID(s): 17969 (mouse), 114553 (rat)

Immunogen

Peptide with sequence C-DEAEDPDPNYAGEP, from the internal region of the protein sequence according to NP_035006.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 48kDa band observed in lysates of cell line Daudi and in preliminary testing of Mouse Spleen lysate (calculated MW of 44.7kDa according to Human NP_000256.4. This molecular weight is routinely observed by other sources. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Expression of the protein seen in the cytoplasm and plasma membrane of NIH3T3 and RAW264.7 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB10589 (1µg/ml) staining of Daudi cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB10589 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and plasma membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB10589 Immunofluorescence analysis of paraformaldehyde fixed RAW264.7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and plasma membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).