

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06343-B - Goat Anti-PAX3, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: PAX3, paired box 3, CDHS, HUP2, WS1, WS3, paired box homeotic

gene 3, paired domain gene 3, paired domain gene HuP2

Official Symbol: PAX3

Accession Number(s): NP\_852122.1; NP\_000429.2; NP\_039230.1; NP\_852123.1;

NP\_852124.1; NP\_852126.1; NP\_852125.1; NP\_001120838.1

Human GenelD(s): 5077

Non-Human GenelD(s): 18505 (mouse), 114502 (rat)

**Important Comments:** This antibody is expected to recognize all 8 reported isoforms

(NP\_852122.1; NP\_000429.2; NP\_039230.1; NP\_852123.1; NP\_852124.1;

NP\_852126.1; NP\_852125.1; NP\_001120838.1).

# **Immunogen**

Peptide with sequence TTLAGAVPRMMC., from the N Terminus of the protein sequence according to NP\_852122.1; NP\_000429.2; NP\_039230.1; NP\_852123.1; NP\_852124.1; NP\_852126.1; NP\_852125.1; NP\_001120838.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 45kDa band observed in Human Duodenum lysates (calculated MW of 45.2kDa according to NP\_852125.1 and 44.8kDa according to NP\_852126.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.3-1µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow

Biotinylated EB06343 (0.5μg/ml) staining of Human Duodenum lysate (35μg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.