



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09134 - Goat Anti-PCSK9 (aa214-228) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PCSK9, proprotein convertase subtilisin/kexin type 9, FH3, HCHOLA3, LDLQCQ1, NARC-1, NARC1, OTTHUMP00000009833, convertase subtilisin/kexin type 9 preproprotein, hypercholesterolemia, autosomal dominant 3, neural apoptosis regulated convertase 1

Official Symbol: PCSK9

Accession Number(s): NP_777596.2

Human GeneID(s): [255738](#)

Immunogen

Peptide with sequence TRFHRQASKCDSHGT, from the internal region of the protein sequence according to NP_777596.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 70-75kDa band observed in HepG2 cell lysates and approx 75kDa in HeLa cell lysates (calculated MW of 74.3kDa according to NP_777596.2)..

Recommended concentration: 1-2g/ml. Primary incubation 1 hour at room temperature.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat

EB09134 (2µg/ml) staining of HeLa (A) and (1ug/ml) of HepG2 (B) cell lysate (35µg protein in RIPA buffer).
Detected by chemiluminescence.

EB09134 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.