

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire

OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05500-T - Goat Anti-Plakoglobin / Gamma-catenin Antibody - Trial

Size: 20µg specific antibody in 40µl



### **Target Protein**

**Principal Names:** JUP, GAMMA-CATENIN, junction plakoglobin, DP3, PDGB, PKGB, DPIII, ARVD12, CTNNG, catenin (cadherin-associated protein), gamma (80kD), catenin (cadherin-associated protein), gamma 80kDa, desmoplakin III, desmoplakin-3

Official Symbol: JUP

Accession Number(s): NP\_002221.1, NP\_068831.1

Human GeneID(s): 3728

Non-Human GenelD(s): 16480 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms, as

represented by NP\_002221.1 and NP\_068831.1.

# Immunogen

Peptide with sequence C-GLRPPYPTADHMLA, from the C Terminus of the protein sequence according to NP\_002221.1, NP\_068831.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 80kDa band observed in lysates of cell lines A431 and NIH3T3 and in Rat Eye lysates (calculated MW of 81.7kDa according to Human NP\_002221.1 and Mouse NP\_034723.1). Recommended concentration: 0.1-0.3µg/ml.

IHC: Paraffin embedded Human Heart. Recommended concentration: 5µg/ml.

**Additional validation:** This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

#### **Species Reactivity**

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

#### Specific Reference

#### This antibody has been successfully used in the following paper:

Krzysztof Sikorski, Adi Mehta, Marit Inngjerdingen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen

A high-throughput pipeline for validation of antibodies

Nat Methods. 2018 Nov;15(11):909-912

PMID: 30377371

EB05500 (0.3μg/ml) staining of A431 (A) and NIH3T3 (B) cell lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB05500 staining (1µg/ml) of Rat Eye lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.

EB05500 (5µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.