

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05234 - Goat Anti-PPP2R5A Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PPP2R5A, protein phosphatase 2, regulatory subunit B (B56), alpha isoform, MGC131915, PP2A, B subunit, B\1 alpha isoform, PP2A, B subunit, B56 alpha isoform, PP2A, B subunit, PR61 alpha isoform, PP2A, B subunit, R5 alpha isoform, serine/threonine protein phosphatase 2A, 56 kDa regulatory subunit, alpha isoform, protein phosphatase 2, regulatory subunit B\1, alpha isoform, B56A, MGC131915, PR61A

**Official Symbol:** PPP2R5A

**Accession Number(s):** NP\_006234.1

**Human GeneID(s):** [5525](#)

### Immunogen

Peptide with sequence C-AYNMHSILSNTSAE, from the C Terminus of the protein sequence according to NP\_006234.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 55kDa band observed in human heart and skeletal muscle lysate (calculated MW of 56.2kDa according to NP\_006234.1). Recommended for use at 0.5-2µg/ml.

**IHC:** Paraffin embedded Human Heart and Brain (Cortex). Recommended concentration: 5µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Rat, Dog, Pig, Cow

EB05234 staining (0.5µg/ml) of human muscle lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

EB05234 (5µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB05234 (5µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.