

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06456 - Goat Anti-PRL1 / PTP4A1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PTP4A1, HH72, PRL1, PRL-1, PTPCAAX1, PTP(CAAX1), protein tyrosine phosphatase type IVA, member 1, Protein tyrosine phosphatase IVA1, RP5-1148A21.2, DKFZp779M0721, protein tyrosine phosphatase type IVA protein 1

Official Symbol: PTP4A1

Accession Number(s): NP_003454.1

Human GeneID(s): [7803](#)

Immunogen

Peptide with sequence CFKDSNGHRNN, from the C Terminus of the protein sequence according to NP_003454.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: This antibody has been successfully used in Western Blot as described in the following papers, PMID: 18997816 and PMID: 36793871.

Immunofluorescence: Strong expression of the protein seen in HeLa and NIH3T3 cells. Recommended concentration: 10µg/ml.

Additional validation: This antibody has been successfully used in the following paper: Achiwa H et al. (2007) PMID: 17234774.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog

Specific References

This antibody has been successfully used in WB on Mouse:

Byungtae Hwang, Min-Gi Kwon, Min Ji Cho, Nam-Kyung Lee, Jangwook Lee, Jeong Woong Lee, Kyoung-Jin Oh, Kwang-Hee Bae, Jung Hwan Hwang, Jeong-Ki Min, Jong-Gil Park

Hepatic PTP4A1 ameliorates high-fat diet-induced hepatosteatosis and hyperglycemia by the activation of the CREBH/FGF21 axis.

Theranostics. 2023 Jan 22;13(3):1076-1090.

PMID: 36793871

This antibody has been successfully used in WB on Human:

Min SH, Kim DM, Heo YS, Kim YI, Kim HM, Kim J, Han YM, Kim IC, Yoo OJ.

New p53 target, phosphatase of regenerating liver 1 (PRL-1) downregulates p53.

Oncogene. 2009 Jan 29;28(4):545-54.

PMID: 18997816

This antibody has been successfully used in the following paper:

Achiwa H, Lazo JS.

PRL-1 tyrosine phosphatase regulates c-Src levels, adherence, and invasion in human lung cancer cells.

Cancer Res. 2007 Jan 15;67(2):643-50.

PMID: 17234774

EB06456 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB06456 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).