

## **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06485 - Goat Anti-SCARF1 / SREC Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** SCARF1, SREC, KIAA0149, MGC47738, scavenger receptor class F, member 1, acetyl LDL receptor, scavenger receptor expressed by endothelial cells **Official Symbol:** SCARF1

Accession Number(s): NP\_003684.2; NP\_663325.1; NP\_663327.2 Human GenelD(s): <u>8578</u>

**Important Comments:** This antibody is expected to recognise all three reported isoforms (NP\_003684.2; NP\_663325.1; NP\_663327.2). Epitope corresponding to aa 20-31, mapping to the mature human SCARF1 / SREC1 protein.

#### Immunogen

Peptide with sequence SELDPKGQHVC, from the internal region of the protein sequence according to NP\_003684.2; NP\_663325.1; NP\_663327.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 85-90kDa band observed in Human Heart lysates (calculated MW of 87.4kDa according to NP\_003684.2). Recommended concentration: 0.3-1.0µg/ml. Primary incubation was 1 hour. An additional faint band of 28kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal.

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human EB06485 (0.3µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Detected by chemiluminescence.