

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05114 - Goat Anti-SHP2 / PTPN11 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PTPN11, SHP2, protein tyrosine phosphatase, non-receptor type 11 (Noonan syndrome 1), CFC, NS1, BPTP3, PTP2C, SHP-2, PTP-1D, SH-PTP2, SH-PTP3, Noonan syndrome 1, protein-tyrosine phosphatase 2C, protein tyrosine phosphatase, non-receptor type 11, MGC14433, protein tyrosine phosphatase-2

Official Symbol: PTPN11

Accession Number(s): NP_002825.3; NP_001317366.1

Human GeneID(s): [5781](#)

Non-Human GeneID(s): 19247 (mouse), 25622 (rat)

Immunogen

Peptide with sequence C-YENVGLMQQQKSFR, from the C Terminus of the protein sequence according to NP_002825.3; NP_001317366.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 70kDa band observed in Human Muscle lysates (calculated MW of 68.0kDa according to NP_002825.3). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the nuclei and cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow

EB05114 staining (2µg/ml) of Human Muscle lysate (RIPA buffer, 35µg total protein per lane). Detected by chemiluminescence.

EB05114 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB05114 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.