

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05904 - Goat Anti-SNX16 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SNX16, sorting nexin 16, DKFZp666H147

Official Symbol: SNX16

Accession Number(s): NP_071416.2; NP_690049.1; NP_690050.1

Human GeneID(s): [64089](#)

Non-Human GeneID(s): 74718 (mouse), 64088 (rat)

Important Comments: This antibody is expected to recognise both isoform a (NP_071416.2; NP_690049.1) and isoform b (NP_690050.1).

Immunogen

Peptide with sequence ATPYVPVPMPIGN-C, from the N Terminus of the protein sequence according to NP_071416.2; NP_690049.1; NP_690050.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


Western blot: Approx 38kDa band observed in lysates of cell line Daudi (calculated MW of 39.2kDa according to NP_071416.2). In transfected HEK293 transiently expressing SNX16 a band of approx. 40kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 0.05-0.2µg/ml.

IHC: Paraffin embedded Human Placenta, Tonsil and Breast. Recommended concentration: 3.75µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow




EB05904 (0.05µg/ml) staining of Daudi lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.




HEK293 overexpressing SNX16 (RC207476) and probed with EB05904 (mock transfection in first lane), tested
by Origene.



EB05904 (3.75µg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05904 (3.75µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05904 (3.75µg/ml) staining of paraffin embedded Human Breast. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.