



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB13088 - Goat Anti-SARS-COV2 Spike protein-S1 Domain (N Terminal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) isolate Wuhan-Hu-1, Spike glycoprotein, E2, Peplomer protein

Official Symbol: SARS-CoV-2

Accession Number(s): P0DTC2; NC_045512.2/MN908947

Immunogen

Peptide with sequence PROPRIETARY.,-C, from the N Terminus of the protein sequence according to P0DTC2; NC_045512.2/MN908947.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 100-110kDa band observed on recombinant SARS-COV2 Spike Protein S1. Recommended concentration 0.5-1ug/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Transient expression of SARS-CoV-2 spike protein expressing plasmid, seen in the cytoplasm of HEK293 transfected cells.

Species Reactivity

Tested: Viral

Expected from sequence similarity: Human Coronavirus

EB13088 (1µg/ml) staining of recombinant Spike Protein S1 (1µg protein in PBS, pH 7.4). Detected by chemiluminescence.

EB13088 Immunofluorescence analysis of paraformaldehyde fixed HEK293 cells transiently transfected with SARS-CoV-2 spike protein expressing plasmid, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: non-transfected HEK293 cells.