

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09721 - Goat Anti-TAF8 / TAFII43 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TAF8, TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 43kDa, 43, FLJ32821, II, TAF, TAFII43, TBN, OTTHUMP00000016392, TAF(II)43, TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 50 kD, TATA box binding protein (TBP)-associated factor, RNA polymerase II, A, 45/50kDa, TBP-associated factor 8, TBP-associated factor TAFII43, TBP-associated factor, RNA polymerase II, 43 kD, taube nuss homolog

Official Symbol: TAF8

Accession Number(s): NP_612639.2

Human GenelD(s): 129685

Non-Human GenelD(s): 63856 (mouse), 316216 (rat)

Immunogen

Peptide with sequence KSYCEHTARTQPT, from the internal region of the protein sequence according to NP_612639.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Preliminary experiments gave an approx 38kDa band in Human Brain (Amygdala, Cerebellum, Hippocampus) lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 34.3kDa according to NP_612639.2. The 38kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

Species Reactivity

Tested:

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow