



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12431 - Goat Anti-TICAM-1 / TRIF Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TICAM1, toll-like receptor adaptor molecule 1, PRVTIRB, TICAM-1, TRIF, TIR domain containing adaptor inducing interferon-beta, TIR domain-containing adapter molecule 1, TIR domain-containing adapter protein inducing IFN-beta, proline-rich, vinculin and TI

Official Symbol: TICAM1

Accession Number(s): NP_891549.1

Human GeneID(s): [148022](#)

Non-Human GeneID(s): 106759 (mouse), 363328 (rat)

Immunogen

Peptide with sequence PDGATFCEDFQVP, from the internal region of the protein sequence according to NP_891549.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 70kDa band observed in lysates of cell line Daudi (calculated MW of 76.4kDa according to NP_891549.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB12431 (2µg/ml) staining of Daudi lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.