

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire

OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11081 - Goat Anti-Transglutaminase 2 (aa578-591) Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** C polypeptide, G-ALPHA-h, GNAH, protein-glutamine gamma-glutamyltransferase 2, protein-glutamine-gamma-glutamyltransferase, TG(C), TG2, TGase C, TGase H, TGase-2, TGase-H, TGC, tissue transglutaminase, transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase),

transglutaminase C, transglutaminase H, transglutaminase-2, TGM2

Official Symbol: TGM2

Accession Number(s): NP\_004604.2

Human GeneID(s): 7052

Non-Human GenelD(s): 21817 (mouse), 56083 (rat)

Important Comments: This antibody is expected to recognize isoform a (NP\_004604.2)

only.

## **Immunogen**

Peptide with sequence C-ERDLYLENPEIKIR, from the internal region of the protein sequence according to NP\_004604.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 80kDa band observed in Mouse and Rat Kidney lysates (calculated MW of 77.3kDa according to Human NP\_004604.2 and 77.0kDa according to Mouse NP\_033399.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow

EB11081 ( $1\mu g/ml$ ) staining of Mouse Kidney lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.