

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06572 - Goat Anti-TRAP2 / Proteasome subunit 26S Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** TRAP2, MGC14274, PSMD2, S2, P97, proteasome (prosome, macropain) 26S subunit, non-ATPase, 2, 55.11 protein, TNFR-associated protein 2, 26S proteasome subunit p97, 26S proteasome regulatory subunit S2, 26S proteasome non-ATPase regulatory subunit 2, tumor necrosis factor receptor-associated protein 2

**Official Symbol:** PSMD2

**Accession Number(s):** NP\_002799.3; NP\_001265637.1; NP\_001265638.1

**Human GeneID(s):** [5708](#)

### Immunogen

Peptide with sequence C-VILRKPNPYDL, from the C Terminus of the protein sequence according to NP\_002799.3; NP\_001265637.1; NP\_001265638.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 100kDa band observed in lysates of cell lines HeLa, U251, NIH3T3 and KNRK (calculated MW of 100kDa according to Human NP\_002799.3, Mouse NP\_598862.1 and Rat NP\_001026809.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

### Species Reactivity

**Tested:** Human, Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow

EB06572 (1 µg/ml) staining of HeLa (A), U251 (B), KNRK (C) and (0.3ug/ml) NIH3T3 (D) cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB06572 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB06572 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.